

# Depression and anxiety are associated with a diagnosis of hypertension 5 years later in a cohort of late middle-aged men and women.

Ginty, Annie; Carroll, Douglas; Roseboom, TJ; Phillips, Anna; de Rooij, SR

DOI:  
[10.1038/jhh.2012.18](https://doi.org/10.1038/jhh.2012.18)

*Document Version*  
Early version, also known as pre-print

*Citation for published version (Harvard):*  
Ginty, A, Carroll, D, Roseboom, TJ, Phillips, A & de Rooij, SR 2013, 'Depression and anxiety are associated with a diagnosis of hypertension 5 years later in a cohort of late middle-aged men and women.', *Journal of Human Hypertension*, vol. 27, pp. 187–190. <https://doi.org/10.1038/jhh.2012.18>

[Link to publication on Research at Birmingham portal](#)

## **Publisher Rights Statement:**

The final published version is available at *Journal of Human Hypertension* (2013) 27, 187–190; doi:10.1038/jhh.2012.18; published online 17 May 2012

## **General rights**

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

## **Take down policy**

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact [UBIRA@lists.bham.ac.uk](mailto:UBIRA@lists.bham.ac.uk) providing details and we will remove access to the work immediately and investigate.

# Depression and anxiety are associated with a diagnosis of hypertension five years later in a cohort of late middle aged men and women.

Running head: Depression, anxiety, and hypertension

Annie T. Ginty, BS<sup>1</sup>, Douglas Carroll, PhD<sup>1</sup>, Tessa J. Roseboom, PhD<sup>2</sup>, Anna C. Phillips, PhD<sup>1</sup>,  
and Susanne R. de Rooij, PhD<sup>2</sup>

<sup>1</sup>School of Sport and Exercise Sciences, University of Birmingham, Birmingham, United Kingdom; <sup>2</sup>Department of Clinical Epidemiology and Biostatistics, Academic Medical Centre, University of Amsterdam, Amsterdam, the Netherlands

Address correspondence to: Susanne R. de Rooij, Department of Clinical Epidemiology and Biostatistics, Academic Medical Centre, University of Amsterdam, Meibergdreef 9, PO Box 22660, 1100 DD Amsterdam, The Netherlands. Tel: \*\*31 20 56 65810; Fax: \*\*31 20 69 12683;  
E-mail: [s.r.derooij@amc.uva.nl](mailto:s.r.derooij@amc.uva.nl)

Word count: 2271

Tables: 2

Figures: 0

## ABSTRACT

**Objective:** To examine the association between symptoms of depression and anxiety and hypertension status. **Methods:** Participants ( $n = 455$ , 238 women) were drawn from the Dutch Famine Birth Cohort Study. In 2002-2004, they attended a clinic assessment during which socio-demographics, anthropometrics, resting systolic blood pressure, and health behaviours were measured. Symptoms of depression and anxiety were measured using the Hospital Anxiety and Depression Scale. In 2008-2009, participants completed a questionnaire which asked whether they ever had a physician diagnosing them as suffering from hypertension. **Results:** In separate regression models that initially adjusted for age and then additionally for sex, socioeconomic status, smoking, sports participation, alcohol consumption, resting systolic blood pressure, antidepressive and anxiolytic medication, whether or not participants were exposed to the Dutch famine *in utero*, BMI, and waist:hip ratio, both depression and anxiety were positively associated with hypertension status. Those who met the criterion for possible clinical depression and anxiety were also more likely to be hypertensive, and these associations remained statistically significant in the fully adjusted regression model. **Conclusion:** Symptoms of depression and anxiety were associated with a diagnosis of hypertension assessed five years later, although the mechanisms underlying these associations remain to be determined.

**Key words:** anxiety, depression, hypertension

## **INTRODUCTION**

A number of recent studies testify to positive cross-sectional and/or prospective associations between major depressive disorder and hypertension (1–3) and between generalised anxiety disorder and hypertension (1). In these studies, depression and anxiety were based on a diagnostic interview. However, evidence of an association between questionnaire assessed symptoms of depression and anxiety and hypertension comes from older studies and is not wholly consistent (4-9). The National Health and Nutrition Examination Survey (4) and the Framingham study (5) observed that high levels of symptoms of depression and anxiety were associated with an increased risk of developing hypertension, whereas the Coronary Artery Risk Development in Young Adults study (6) found no consistent relationship between symptoms of depression and anxiety and hypertension. Other studies have demonstrated associations between depression and anxiety symptomatology and lower blood pressure (7-9). Accordingly, we judged it timely to re-visit this issue and data collected as part of the Dutch Famine Birth Cohort Study (10) permitted re-examination of the association between symptoms of depression and anxiety and hypertension status. We hypothesised, on the balance of the previous evidence, that elevated depression and anxiety symptom levels would be associated with an increased risk for hypertension.

## **METHOD**

### **Participants**

Participants were selected from the Dutch Famine Birth Cohort, which comprises men and women who were born in Amsterdam, the Netherlands, between November 1943 and February 1947. The selection procedures and subsequent loss to follow up have been described in detail

elsewhere (11-13). In 2002-2004, 740 members of the cohort provided data at a clinic assessment, performed by a trained research nurse. In 2008-2009, 455 of those who had attended the clinic returned a completed questionnaire package, which among other things, asked about diagnosed illness. The study was approved by the local Medical Ethics Committee, carried out in accordance with the Declaration of Helsinki, and with the adequate understanding and written consent of the participants.

## **Measures**

In 2002-2004, socio-demographics, life style, and anthropometrics were assessed. Socio-economic status (SES) was measured using the ISEI (International Socio-Economic Index)-92, which is based on the participant's or their partner's occupation, whichever has the higher status (14). Measured values in the ISEI-92 scale ranged from 16 (low status, for example a cleaning person) to 87 (high status, for example a lawyer). Participants were asked whether they were current smokers, ex-smokers, or had never smoked. They were also asked whether they participated in sports and how many alcoholic drinks of different types they consumed per week, and an overall units per week measure was derived. Participants had to indicate whether they were taking antidepressant and/or anxiolytic medication (NSRIs). Height was measured twice using a fixed or portable stadiometer and weight twice using Seca and portable Tefal scales. Body mass index (BMI) was computed in  $\text{kg/m}^2$  from the averages of the two height and weight measurements. Waist and hip circumference was measured in duplicate with a flexible tape measure and an average taken. Waist was measured midway between the costal margin and the iliac crest and hip at the widest point over the buttocks to compute the waist:hip ratio. Systolic blood pressure (SBP) was measured twice in a row on two occasions (morning and afternoon)

using an automated device (Omron 705 CP/IT; Omron Healthcare UK, West Sussex, UK) and appropriate cuff sizes. Mean SBP was calculated using all available measurements.

Symptoms of depression and anxiety were measured using the Hospital Anxiety and Depression Scale (HADS) (15). The HADS is a well-recognised assessment instrument that comprises 14 items, seven measuring depression and seven measuring anxiety. The depression subscale emphasises anhedonia and excludes somatic items. Items are scored on a 4-point scale, 0 to 3; the higher the score, the greater the depression and anxiety. The HADS has good concurrent validity (16,17), performs well as a psychiatric screening device (17,18), and has been shown to have acceptable psychometric properties; for example, a Cronbach's  $\alpha$  of .90 for the depression items and .93 for the anxiety items has been reported (19) and test-retest reliability coefficients as high as .85 for depression and .84 for anxiety have been found (17). Possible caseness for both depression and anxiety is defined by scores  $\geq 8$ .

The 2008-2009 questionnaire asked participants whether or not they had ever received a diagnosis of hypertension from a physician. The mean (SD) temporal lag between this assessment and the measurement of depression and anxiety was 5.5 (0.6) years.

## **Statistical Analyses**

Preliminary analyses of differences between participants with and without a diagnosis of hypertension were conducted using  $\chi^2$  and t-tests. The main analysis was by logistic regression, where depression and anxiety served as the independent variables in separate models with hypertension status as the dependent variable. The initial models adjusted only for age at clinic

assessment, whereas subsequent models additionally adjusted for sex, SES, smoking, sports participation, alcohol consumption, SBP, antidepressive and anxiolytic medication, whether or not participants were exposed to the Dutch famine *in utero*, BMI, and waist:hip ratio. These covariates were chosen because they have been implicated in depression, anxiety, and/or hypertension. Depression and anxiety were first tested as continuous variables, i.e. actual HADS scores, and subsequently as categorical variables, i.e. HADS caseness.

## RESULTS

The summary characteristics of the cohort by hypertension status are presented in Table 1. Two hundred and nineteen (48%) of the participants reported a diagnosis of hypertension at the second time point of assessment. HADS depression and anxiety scores were highly correlated,  $r(433) = .68, p < .001$ , as was HADS depression and anxiety caseness,  $C(\text{coefficient of contingency})(433) = .43, p < .001$ . In regression models adjusting only for age, both depression,  $OR = 1.13, 95\%CI\ 1.05 - 1.21, p = .001$ , and anxiety scores,  $OR = 1.11, 95\%CI\ 1.05 - 1.18, p = .001$ , were associated with future hypertension status; the greater the symptoms of depression and anxiety the greater the likelihood that the particular person would have been diagnosed with hypertension five years later. These associations remained statistically significant in the fully adjusted models: for symptoms of depression,  $OR = 1.19, 95\%CI\ 1.09 - 1.30, p < .001$ , and for symptoms of anxiety,  $OR = 1.14, 95\%CI\ 1.06 - 1.23, p = .001$ . In these models, hypertension was also associated with a higher BMI and a higher resting SBP at the clinic assessment,  $p = .001$  in both cases. Men were also more likely to be hypertensive in the depression model,  $p = .05$ .

These analyses were repeated using cut-off scores of  $\geq 8$  to identify possible clinical depression and anxiety. In the age adjusted models, both depression caseness, OR = 2.34, 95%CI 1.14 – 4.79,  $p = .02$ , and anxiety caseness, OR = 1.73, 95%CI 1.08 – 2.77,  $p = .02$ , were positively associated with hypertension status. With full adjustment, the association between depression caseness and hypertension remained significant, OR = 3.44, 95%CI 1.44 – 8.23,  $p = .005$ , as did the association between anxiety caseness and hypertension, OR = 2.08, 95%CI 1.17 – 3.72,  $p = .01$ .

### **Sensitivity analyses**

During the clinic assessment, 108 participants admitted to taking antihypertensive medication. Accordingly, it could be argued that the present prospective associations between symptoms of depression and anxiety, on the one hand, and hypertension, on the other, were largely reflecting cross-sectional associations. As a consequence we first repeated the analyses for the continuous depression and anxiety scores removing those individuals from the analyses. In the fully adjusted models, both symptoms of depression, OR = 1.18, 95%CI 1.06 – 1.31,  $p = .003$ , and anxiety, OR = 1.14, 95%CI 1.03 – 1.25,  $p = .008$ , continued to predict hypertensive status five years later. We also repeated the analyses of the whole sample with hypertensive medication status added as a further covariate to the fully adjusted models. Although the associations were slightly attenuated, symptoms of depression, OR = 1.16, 95%CI 1.06 – 1.28,  $p = .002$ , and symptoms of anxiety, OR = 1.11, 95%CI 1.02 – 1.21,  $p = .01$ , continued to predict future hypertension.



## DISCUSSION

Almost half the current sample reported that a physician had diagnosed them as hypertensive. This is a higher prevalence than that reported in other studies (1,3), but could reflect the age of the sample who were all over 60 at the time of the questionnaire assessment of hypertension status as well as the close medical attention paid to this cohort. It certainly suggests, though, that there was little undiagnosed hypertension, a concern when relying on self-reports of physician diagnosis (1). As hypothesised, participants with high levels of depression and anxiety symptoms were more likely to be hypertensive when questioned five years later. Hypertensive participants were also more likely to meet the criterion for possible clinical caseness for depression and anxiety at the earlier assessment. Overall, our results are in line with those from other studies comparing hypertension in those varying in symptoms of depression and anxiety (4,5) and those with and without Major Depressive Disorder (1-3) and with and without Generalised Anxiety Disorder (1).

There are several possible pathways through which depression and anxiety might contribute to the development of hypertension (20). The two most cited are unhealthy behaviours and physiological dysregulation. However, in the present analyses, the associations between symptoms of depression and anxiety and hypertension were still evident after adjustment for three of the more prominent unhealthy behaviours, smoking, non participation in sports and high levels of alcohol consumption. With regard to physiological dysregulation, altered activity of the hypothalamic-pituitary-adrenal axis has been observed in approximately 50% of depressed

patients (21), and this, in turn, may increase the risk of hypertension. Cortisol was measured in the present study during the clinic assessment, but the addition of basal cortisol to our regression models left the outcomes unaltered. Altered autonomic function have also been suggested as a possible psychophysiological mechanism (22). However, in a previous study in the same cohort we found that those with symptoms of depression and anxiety had decreased and not increased cardiovascular stress responses (23). In the present study, the addition of SBP and HR reactivity to our regression models left the outcomes unaltered ( $p < .001$  and  $p = .001$  for depression and anxiety, respectively). One other possibility is reverse causation, such that increased symptoms of depression and anxiety are reactions to a diagnosis of hypertension, a chronic and ultimately life-threatening condition. However, when we excluded participants who reported taking antihypertensive medication at the time of the assessment of depression and anxiety, symptoms of depression and anxiety continued to predict hypertension status at the subsequent assessment, making reverse causation an unlikely explanation.

The present study is not without limitations. First, from observational data it is impossible to determine causality and the direction of causality. However, as indicated above, reverse causation would seem to be an unlikely explanation. Nevertheless, the issue of confounding is ever present in observational studies (24). However, we did adjust statistically for a broad range of potential confounders, including whether or not participants were exposed to the Dutch famine and SBP during the clinic assessment, and, in sensitivity analyses, whether or not participants reported taking antihypertensive medication at the earlier assessment. That the links between depression and anxiety and future hypertension survived such adjustments suggest that these associations were not simply an artefact of cross-sectional relationships. Second, we relied on a

questionnaire measure of symptoms rather than formal diagnoses of major depressive and anxiety disorders. Further, use of the HADS may have lead to an underestimation of the association between depression and hypertension as it excludes somatic-affective symptoms of depression and anxiety and these may particularly increase the risk for hypertension. However, our findings are very much in accord with those from studies that did examine hypertension in the context of psychiatric diagnoses (1-3). Third, the relative seniority of our participants and the consequently high prevalence of diagnosed hypertension could be regarded as limiting the generalisability of our findings. However, given that similar associations have emerged in middle-aged cohorts (1-3), it would appear that the relationship between depression and anxiety and hypertension transcends the age of the sample studied. Fourth, this study relied on self-report of a physician diagnosis of hypertension. However, the high prevalence of hypertension suggests that there were unlikely to be many false negatives. Further, other studies in the field have also used self-report of hypertension (2,3). Finally, although the high prevalence of hypertension makes it unlikely, it cannot be ruled out that depressed and anxious subjects were more likely to be diagnosed with hypertension and non-depressed / anxious subjects remained undiagnosed, as hypertension is frequently asymptomatic and depressed / anxious subjects may have been checked more often. Another possibility would be that there were more false positives among depressed /anxious subjects due to panic-attack related high blood pressure being misinterpreted as hypertension.

In conclusion, the present analyses indicated that symptoms of depression and anxiety were associated with a diagnosis of hypertension assessed five years later. These associations would not seem to be accounted for by unhealthy behaviour, nor would they appear a function of

reverse causation, where increased symptoms of depression and anxiety are reactions to a diagnosis of hypertension. Further studies are needed to unravel the mechanisms underlying what appear to be fairly robust associations between psychological status and hypertension.

## **Acknowledgements**

We thank the participants for their willing cooperation. The present study was supported by the Netherlands Heart Foundation NHS2007B083, the European Science Foundation (EuroSTRESS) and the Netherlands Organisations for Scientific Research (NWO). These funding sources had no role in study design, data collection, analysis and interpretation of the data.

## **Conflict of interest**

The authors declare no conflict of interest.

## References

1. Carroll D, Phillips AC, Gale CR, Batty D. Generalized anxiety and major depressive disorders, their comorbidity and hypertension in middle-aged men. *Psychosom Med* 2010; **72**: 16-19.
2. Patten SB, Beck CA, Kassam A, Williams JVA, Barbui C, Metz LM. Long-term medical conditions and major depression: strength of association in the general population. *Can J Psychiatry* 2005; **50**: 195-202.
3. Patten SB, Williams JVA, Lavorato DH, Campbell NRC, Eliasziw M, Campbell TS. Major depression as a risk factor for high blood pressure: Epidemiologic evidence from a national longitudinal study. *Psychosom Med* 2009; **71**: 273-279.
4. Jonas BS, Franks P, Ingram DD. Are symptoms of anxiety and depression risk factors for hypertension? Longitudinal evidence from the National Health and Nutrition Examination Survey I Epidemiologic Follow-up Study. *Arch Fam Med* 1997; **6**: 43-49.
5. Markovitz JH, Matthews KA, Kannel WB, Cobb JL, D'Agostino RB. Psychological predictors of hypertension in the Framingham Study. Is there tension in hypertension? *JAMA* 1993; **270**: 2439-2443.
6. Yan LL, Liu K, Matthews KA, Daviglus ML, Ferguson TF, Kiefe CI. Psychosocial factors and risk of hypertension: the Coronary Artery Risk Development in Young Adults (CARDIA) study. *JAMA* 2003; **290**: 2138-2148.
7. Hildrum B, Mykletun A, Stordal E, Bjelland I, Dahl AA, Holmen J. Association of low blood pressure with anxiety and depression: the Nord-Trøndelag Health Study. *J Epidemiol Community Health* 2007; **61**: 53-58.
8. Stroup-Benham CA, Markides KS, Black SA, Goodwin JS. Relationship between low blood pressure and depressive symptomatology in older people. *J Am Geriatr Soc* 2000; **48**: 250-255.
9. Barrett-Connor E, Palinkas LA. Low blood pressure and depression in older men: a population based study. *BMJ* 1994; **308**: 446-449.
10. Roseboom T, de Rooij S, Painter R. The Dutch famine and its long-term consequences for adult health. *Early Hum Dev* 2006; **82**: 485-491.
11. Ravelli AC, van der Meulen JH, Michels RP, Osmond C, Barker DJ, Hales CN, Bleker OP. Glucose tolerance in adults after prenatal exposure to famine. *Lancet* 1998; **351**: 173-177.

12. Painter RC, Roseboom TJ, Bossuyt PM, Osmond C, Barker DJ, Bleker OP. Adult mortality at age 57 after prenatal exposure to the Dutch famine. *Eur J Epidemiol* 2005; **20**: 673-676.
13. de Rooij SR, Veenendaal MV, Raikkonen K, Roseboom TJ. Personality and stress appraisal in adults prenatally exposed to the Dutch famine. *Early Hum Dev* 2011; epub ahead of print, Sep 26.
14. Bakker B, Sieben I. Maten voor prestige, social-economische status en sociale klasse voor de standard beroepenclassificatie 1992. *Sociale Wetenschappen* 1997; **40**: 1-22.
15. Zigmond AS, Snaith RP. The Hospital Anxiety and Depression Scale. *Acta Psychiatr Scand* 1983; **67**: 361-370.
16. Bramley PN, Easton AME, Morley S, Snaith RP. The differentiation of anxiety and depression by rating scales. *Acta Psychiatr Scand* 1988; **77**: 133-138.
17. Herrmann, C. International experiences with the Hospital Anxiety and Depression Scale- a review of validation data and clinical results. *J Psychosom Res* 1997; **42**: 17-41.
18. Bjelland I, Dahl AA, Haug TT, Neckelmann D. The validity of the Hospital Anxiety and Depression Scale: an updated literature review. *J Psychosom Res* 2002; **52**: 69-77.
19. Moorey S, Greer S, Watson M, Gorman C, Rowden L, Tunmore R, Robertson B, Bliss J. The factor structure and factor stability of the Hospital Anxiety and Depression Scale in patients with cancer. *Br J Psychiatry* 1991; **158**: 255-259.
20. Beevers, DG, MacGregor, GA. *Hypertension in Practice*. Dunitz: London. 1995.
21. Brown ES, Varghese FP, McEwen BS. Association of depression with medical illness: does cortisol play a role? *Biol Psychiatry* 2004; **55**: 1-9.
22. Kibler JL, Ma M. Depressive symptoms and cardiovascular reactivity to laboratory behavioral stress. *Int J Behav Med* 2004;**11**: 81-87.
23. de Rooij SR, Schene AH, Phillips DI, Roseboom TJ. Depression and anxiety: Associations with biological and perceived stress reactivity to a psychological stress protocol in a middle-aged population. *Psychoneuroendocrinology* 2010;**35**: 866-877.
24. Christenfeld, NJ, Sloan RP, Carroll D, Greenland S. Risk factors, confounding and the illusion of statistical control. *Psychosom Med* 2004; **66**: 737-743.

Table 1. Descriptive characteristics of those with and without a diagnosis of hypertension.

	Hypertensive ( <i>n</i> = 219)		Normotensive ( <i>n</i> = 236)		<i>P</i>
	Mean	SD	Mean	SD	
Age at clinic assessment (years)	58.16	0.88	58.28	0.94	.16
Socio-economic status (ISEI-92)	49.82	13.88	50.88	13.79	.42
Body mass index (kg/m <sup>2</sup> )	30.42	5.04	27.08	3.94	<.001
Waist:hip ratio	93.69	9.00	91.89	8.58	.03
Systolic blood pressure (mmHg)	142.83	16.17	128.68	14.76	<.001
Alcohol consumption (units/week)	9.54	16.53	9.97	12.56	.76
HADS depression score	3.66	3.44	2.64	2.43	<.001
HADS anxiety score	5.99	3.55	4.85	2.97	<.001
	<i>n</i> (%)		<i>n</i> (%)		<i>P</i>



Sex			.23
Male	98 (45)	119 (55)	
Female	121 (51)	117 (49)	
Smoking			.13
Current smokers	44 (42)	61 (58)	
Ex-smokers	83 (46)	97 (54)	
Never smoker	91 (54)	78 (46)	
Sports participation	129 (59)	138 (58)	.93
Exposed to famine <i>in utero</i>	90 (41)	99 (42)	.22
Medication			
Antidepressive	14 (6)	12 (5)	.55
Anxiolytic	18 (8)	18 (8)	.90
Caseness			
Depression	25 (12)	12 (5)	.01
Anxiety	54 (26)	37 (16)	.02

---

Table 2. Summary Table

---

*What is known about this topic*

- Recent studies consistently indicate that major depressive disorder and generalised anxiety disorder are positively related to hypertension.
- Evidence linking symptoms of depression and anxiety with hypertension is less compelling.

*What this study adds*

- Participants with high levels of depression and anxiety symptoms are more likely to be hypertensive five years later.
  - These associations would not seem to be accounted for by unhealthy behaviours.
  - They would also not appear to be a function of reverse causation, where increased symptoms of depression and anxiety are reactions to a diagnosis of hypertension.
-